Producing sons reduces lifetime reproductive success of subsequent offspring in pre-industrial Finns

Ian J. Rickard^{*}, Andrew F. Russell and Virpi Lummaa

Department of Animal and Plant Sciences, University of Sheffield, Sheffield S10 2TN, UK

Life-history theory states that reproductive events confer costs upon mothers. Many studies have shown that reproduction causes a decline in maternal condition, survival or success in subsequent reproductive events. However, little attention has been given to the prospect of reproductive costs being passed onto subsequent offspring, despite the fact that parental fitness is a function of the reproductive success of progeny. Here we use pedigree data from a pre-industrial human population to compare offspring life-history traits and lifetime reproductive success (LRS) according to the cost incurred by each individual's mother in the previous reproductive cost, we hypothesize that individuals born to mothers who previously produced sons will display compromised survival and/or LRS, when compared with those produced following daughters. Controlling for confounding factors such as socio-economic status and ecological conditions, we show that those offspring born after elder brothers have similar survival but lower LRS compared with those born after elder sisters. Our results demonstrate a maternal cost of reproduction manifested in reduced LRS of subsequent offspring. To our knowledge, this is the first time such a long-term intergenerational cost has been shown in a mammal species.

Keywords: evolutionary ecology; intergenerational cost of reproduction; lifetime reproductive success; Lack's hypothesis; Trivers–Willard hypothesis

1. INTRODUCTION

Life-history theory states that reproductive events confer costs upon parents (Roff 1992). While studies of wild populations of animals have shown that reproduction is associated with reduced survival, attractiveness and/or future reproductive capacity for breeders, such evidence is by no means universal (Clutton-Brock 1991). Such a lack of evidence for reproductive costs is commonly attributed to high-quality individuals or those breeding in favourable circumstances being 'immune' to reproductive costs. However, because a parent's fitness depends not only on its own lifetime fecundity, but also that of its offspring, costs may be inter- as well as intragenerational, particularly in species that are expected to favour their own residual reproductive value at the expense of their offspring (Drent & Daan 1980).

In mammals as well as birds, several studies have tested the hypothesis (Lack 1947, 1948*a*, *b*) that the growth and survival of offspring is compromised in those whose parents provisioned artificially enlarged broods or litters, with equivocal results (Hare & Murie 1992; Mappes *et al.* 1995; Kunkele 2000). However, most studies that have tracked the effects of decreased offspring investment on the success of those offspring have only done so as far as independence or recruitment age. The intergenerational cost of reproduction will remain elusive to such short-term observation if the costs of reduced parental investment are

Electronic supplementary material is available at http://dx.doi.org/10. 1098/rspb.2007.1051 or via http://www.journals.royalsoc.ac.uk.

not apparent until the offspring's reproductive life. Only a few studies have investigated parental costs in terms of the reproductive performance of their offspring (Gustafsson & Sutherland 1988; Blondel *et al.* 1998; Koskela 1998; Naguib *et al.* 2006). Perhaps due largely to temporal and logistic constraints of field studies or to the problem of natal dispersal, none has investigated the costs in terms of offspring's lifetime reproductive success.

In this study, we investigate for the first time intergenerational costs of reproduction, measured in terms of offspring's lifetime reproductive success (LRS). We avoid the above logistic problems by using a large pedigree dataset of pre-industrial Finns, which allows following the reproductive events of all offspring throughout their life (see §2). We focus on a differential in cost between individual reproductive events in females and follow the consequent survival and LRS of their subsequent offspring. The differential reproductive cost we use is that of producing male over female offspring. In several mammal species, including humans, there is clear evidence that producing sons is more costly to mothers than producing daughters (table 1). While evidence for a higher cost of males is not ubiquitous (Hewison & Galliard 1999), this has so far manifested itself in those producing sons as greater parasitic load (Festa-Bianchet 1989), delay in return to oestrus (Hogg et al. 1992), reduced probability of future reproduction (Clutton-Brock et al. 1981), reduced post-partum weight accumulation (Birgersson 1998) and reduced maternal longevity (Helle et al. 2002). In addition, in both bighorn sheep, Ovis canadensis (Bérubé et al. 1996), and humans

^{*}Author for correspondence (i.rickard@sheffield.ac.uk).

species	nature of cost demonstrated	study	
bighorn sheep, Ovis canadensis	ewes had higher faecal lungworm count after producing a son	Festa-Bianchet (1989)	
	ewes experienced a delay in return to oestrus after producing a son	Hogg et al. (1992)	
	ewes more likely to have a daughter after producing a son	Bérubé et al. (1996)	
	ewes produced lighter offspring after producing a son	Bérubé et al. (1996)	
red deer, Cervus elaphus	hinds less likely to reproduce in the year following a son; delayed calving in those who did	Clutton-Brock et al. (1981)	
	lower post-reproductive maternal survival after weaning a son (subordinate hinds only)	Gomendio et al. (1990)	
	milk composition dependent on sex, e.g. higher protein percentage in milk provisioned for sons	Landete-Castillejos et al. (2004)	
fallow deer, Dama dama	lower maternal weight accumulation during the period after gestation of offspring, when a son is produced (old hinds only)	Birgersson (1998)	
humans	birth interval longer after the birth of a son (low-parity women only)	Mace & Sear (1997)	
	maternal longevity associated negatively with the number of sons produced	Helle et al. (2002)	
	lower birth weight of offspring produced after sons	Trotnow <i>et al.</i> (1976); Magnus <i>et al.</i> (1985); Blanchard & Ellis (2001); Côté <i>et al.</i> (2003)	
	higher maternal energy intake in women carrying sons	Tamimi et al. (2003)	

(Trotnow *et al.* 1976; Magnus *et al.* 1985; Blanchard & Ellis 2001; Côté *et al.* 2003), the offspring born immediately following a son may have lower birth weight than those born after a female. Although birth weight itself may be only a weak predictor of reproductive success in the bighorn sheep (Festa-Bianchet *et al.* 2000), in humans it has long been known to be associated with early survival (Karn & Penrose 1952) and health in adulthood (Barker 1998) and has also been linked to reproductive success among men (Phillips *et al.* 2001; Vågerö & Modin 2002). Thus, there exists in humans a clear precedent for the development of offspring to be affected by the sex of the offspring an individual's mother has produced in the previous reproductive event.

Consequently, we investigate the relationship between the sex of the offspring produced in one reproductive episode and the survival and LRS of offspring produced immediately after, using 521 mothers (hereafter referred to as P_1) and 1765 of their male and female offspring (hereafter referred to as F_1) born during the eighteenth and nineteenth centuries in Finland. We examined whether elder sibling sex affected in F_1 individuals: (i) the probability of surviving to adulthood and (ii) LRS. In addition, we examined (iii) the mechanism through which elder sibling sex affected LRS, namely through lifetime fecundity and/or survival of offspring (F_2).

2. MATERIAL AND METHODS

(a) Demographic data

To investigate how producing sons affects the success of the following offspring in humans, we collected three generations of pedigree data from historical population registers of The Lutheran Church in Finland. Since the seventeenth century, records have been taken of all births, marriages, deaths and inter-parish movements in Finland, making these one of the most reliable sources of demographic data available for a pre-industrial human population. Social class, an important potential confounding factor in studies concerning the allocation of resources between reproduction and selfmaintenance (van Noordwijk & Dejong 1986) or production of sons and daughters (Trivers & Willard 1973), is based on the occupation of a husband in the family, which is a good correlate of resources available (Karskela 2001). We categorized each family as poor (e.g. farmless families, servants), middle class (e.g. tenant farmers, smiths, sailors) or wealthy (e.g. priests, officers, farm owners, shipmasters; for details see Pettay *et al.* 2007).

We initially followed the survival and reproductive events of a total of 653 reproductive women (the P generation) from five farming/fishing parishes (Lummaa *et al.* 2007; Pettay *et al.* 2007) born during the years 1709–1815 and recorded full life-history data for their offspring born (the F_1 generation), as well as birth and subadult mortality data of the F_2 generation. Here we investigate how the sex of the elder sibling affected the survival and LRS of the F_1 generation.

All F₁ individuals were born between the years 1734 and 1861. Consequently, the study period ended before the availability of reliable contraception, freely available healthcare and the associated transition to low mortality and fertility in Finland, which was not complete until the mid-twentieth century (Korpelainen 2003). The mating system was monogamous to an unusually high degree, with 99% of reproductive individuals in our sample being married. Divorce was forbidden, and so remarriage was permitted only in the event of spousal death. Subadult mortality was high, with only 61% of F₁ individuals in our sample surviving to age 15 (the youngest age of first reproduction recorded in this population). For analysis, we removed cases that did not meet the requirements needed to address the questions of this study (table 2). These were those who were firstborn to their mothers, those who were twins and born to mothers who

Table 2. Sequence of criteria for the inclusion of F_1 data and corresponding sample sizes.

criteria for inclusion	no.	use in analysis
all F ₁ generation	4515	
not firstborn	3811	
not a twin	3315	
elder sibling not a twin	3100	
survival to age 15 known	2769	
lifespan of elder sibling known	1862	
elder sibling sex known	1854	
data available for potential confounders	1842	
birth order < 11	1765	core data: analysis (i)
survived to age 15	1081	
tracked for reproductive lifespan	716	
social class known	656	subset 1: analyses (ii), LRS and (iii), fecundity
produced at least one child	557	subset 2: analysis (iii), offspring survival rate

previously produced twins. We also removed those whose mother had produced more than nine elder siblings previously (4% of remaining cases), because family size is highly skewed and the response terms have high degrees of variance in those of late birth order which decrease modelfitting power. Removal of these cases, plus those for which data were missing, provided a core of 1765 individuals from 521 mothers. Table 3 provides the descriptives relating to the number of individuals in the core data according to their own sex and the sex of their mother's previous offspring.

(b) Statistical methods

Statistical analyses used to address each of the following three questions were conducted using SAS (SAS Institute Inc., release v. 9.1, 2002-2003). In all analyses (§2b(i)-(iii)), parish, year of birth, social class (rich, middle and poor), birth order, interval separating the birth of the focal individual from that of their elder sibling, the total family size (number of F1 individuals surviving to age 15) and own sex were entered sequentially into the analyses to control for potential confounding sources of variation. Those confounding terms that were significant were retained and those that were not were dropped from the model. Once the minimal model was found, the sex of an individual's elder sibling (our term of interest) was added to the final model and its significance determined. All two- and three-way interactions involving elder sibling sex were then tested, but none were significant. Finally, the identity of P mothers was fitted as a random term to account for the use of repeated offspring within families. Generalized linear mixed models (GLMMs) were conducted using the GENMOD function in SAS (SAS 1990). All p-values are two-tailed and significance levels are set at 0.05.

(i) Elder sibling sex and the probability of surviving to age 15

Survival to age 15 (0/1) was considered as a binary response term in a GLMM with logit link function and binomial denominator fixed at 1. After controlling for confounding terms (see \$2b), we tested the influence of elder sibling sex on the survival probability of all individuals in the core data, 1765 F₁ offspring delivered by 521 P mothers (1–9 measures per mother). Table 3. Number of individuals according to own sex and elder sibling sex in the core data, subset 1 and subset 2.

		elder sibling female	elder sibling male	total
core data	focal individual female	407	454	861
	focal individual male	401	503	904
	total	808	957	1765
subset 1	focal individual female	141	161	302
	focal individual male	171	183	354
	total	312	344	656
subset 2	focal individual female	124	137	261
	focal individual male	146	150	296
	total	270	287	557

(ii) Elder sibling sex and LRS

LRS was measured as the lifetime number of F₂ offspring raised to 15 years. This was considered as a continuous response term in a GLMM with Poisson error structure and logarithm link function. Controlling for confounding terms, we tested the influence of elder sibling sex on LRS in a subset of the core data, comprising all those who survived to age 15, who had been followed for their entire reproductive life and those whose social class (an important confounding factor) was known. The end of reproductive lifespan was determined for women as the estimated age of menopause (45 years) and for men as the age by which 90% of men in the whole population had finished reproducing (50 years). This subset comprised 656 individuals, delivered by 322 mothers (1-7 measures per mother). Tables 3 and 4 provide the descriptives of those in this data subset. Since LRS can potentially be negatively influenced by failure to successfully follow all offspring to age 15, the response term was weighted by the proportion of an individual's children whose survivorship to this age was known. Childlessness in this subset (99 individuals, 15%) was associated with the failure to marry among those dying prematurely. Of those who failed to reproduce, less than half (47 individuals, 47%) were married, whereas among those who fathered or gave birth to at least one child, 99% were married.

(iii) Elder sibling sex and mechanisms of LRS

Two life-history traits will govern our measure of LRS: (i) lifetime fecundity and (ii) the survivorship of offspring to age 15. Lifetime fecundity was considered as a continuous response term in a GLMM with Poisson error structure and logarithm link function. Controlling for confounding terms, the influence of elder sibling sex on lifetime fecundity was analysed using the same subset of data as used in analysis (\$2b(ii)).

The proportion of offspring surviving to age 15 was examined by considering LRS as a response term in a GLMM with logit link function and a variable binomial denominator equal to fecundity. After controlling for confounding terms, we tested the influence of elder sibling sex on the survival probability of a F_1 individual's offspring using a further subset of the data. This was because only those individuals with lifetime fecundity of at least 1 could be included in this analysis. This second subset comprised 557 individuals, delivered by 298 mothers (1–7 measures per mother). Tables 3 and 4 provide the descriptives of those in this data subset. As with analysis (§2*b*(ii)), the response term was weighted by

Table 4. Descriptive statistics of reproductive parameters of those F_1 offspring who survived to age 15 themselves and	l were
successfully followed until the end of potential reproductive life (data subset 1), split by sex and social class (all mean ±	s.d.).
(Lifespan here thus refers to the total number of years lived by those who survived to adulthood (age 15).)	

social class	sex	total N	lifespan	age at marriage	age at first reproduction	age at last reproduction	lifetime fecundity	number of sur- viving offspring
rich	m	178	54.82 ± 16.30	26.83 ± 5.40	27.85 ± 5.45	39.78 ± 8.32	4.81 ± 3.26	2.47 ± 2.11
	f	140	61.03 ± 16.27	24.73 ± 5.09	25.52 ± 4.59	38.37 ± 5.10	5.09 ± 3.05	2.94 ± 2.26
middle	m	118	58.82 ± 16.42	27.62 ± 5.11	29.06 ± 6.98	41.44 ± 8.27	4.81 ± 3.06	2.66 ± 2.39
	f	108	63.76 ± 16.97	27.11 ± 6.30	27.27 ± 5.69	38.60 ± 6.20	4.51 ± 2.99	2.48 ± 1.94
poor	m	58	48.42 ± 18.42	27.21 ± 5.40	27.48 ± 5.11	37.95 ± 8.88	2.41 ± 3.02	1.17 ± 1.63
1	f	54	52.50 ± 19.34	28.45 ± 6.94	28.52 ± 5.70	35.40 ± 6.13	2.11 ± 2.11	0.96 ± 1.37
all	m	354	54.41 ± 16.89	27.15 ± 5.29	28.25 ± 6.03	40.19 ± 8.40	4.42 ± 3.27	2.32 ± 2.19
	f	302	60.50 ± 17.49	26.17 ± 5.98	26.63 ± 5.30	38.01 ± 5.77	4.35 ± 3.07	2.42 ± 2.13

the proportion of an individual's children whose survivorship to age 15 was known.

3. RESULTS

(a) Elder sibling sex and the probability of surviving to age 15

Sixty-one per cent of the offspring in the core data survived to age 15. Survival differed between parishes $(\chi_4^2=36.09, p<0.0001)$ and was negatively associated with birth order $(\chi_1^2=3.75, p=0.053)$ and positively with family size $(\chi_1^2=84.67, p<0.001)$. Those born after a long birth interval $(\chi_1^2=4.46, p=0.035)$ and those born later in the time series $(\chi_1^2=3.44, p=0.064)$ were also more likely to survive. After controlling for these effects, we found no evidence to suggest that survival to age 15 was associated with elder sibling sex $(\chi_1^2=0.00, p=0.98;$ figure 1). Nonsignificant potential confounding terms were (mother's) social class $(\chi_2^2=0.81, p=0.67)$ and own sex $(\chi_1^2=0.16, p=0.69)$. Owing to collinearity, family size but not birth order was included in the final model.

(b) Elder sibling sex and LRS

In the subset of data comprising those offspring who survived to age 15 themselves and were successfully followed until the end of potential reproductive life, the median number of children raised to age 15 was 2 and the maximum was 9. A large proportion of individuals (27%) raised no children to adulthood and thus had an LRS of zero. LRS differed marginally between parishes ($\chi_4^2 = 8.79$, p=0.067) and was higher for those in the rich and middle social classes than those in the poor class ($\chi^2_2 = 47.18$, p < 0.0001, adjusted means \pm s.e. 2.83 ± 0.15 , 2.75 ± 0.17 and 1.09 ± 0.15 for rich, middle class and poor, respectively). After controlling for these effects, we found that LRS was significantly higher (27%) in those born after an elder sister than in those born after an elder brother $(\chi_1^2 = 12.01, p = 0.0005;$ figure 2). Finally, LRS was not significantly associated with year of birth ($\chi_1^2 = 0.28$, p=0.60), birth order ($\chi_1^2=0.49$, p=0.48), family size ($\chi_1^2=0.43$, p=0.51), birth interval ($\chi_1^2=0.01$, p=0.92) or own sex ($\chi_1^2=0.30$, p>0.58).

(c) Elder sibling sex and mechanisms of LRS

Among those who survived to age 15 and were successfully followed until the end of potential reproductive life, the median lifetime fecundity was 4 and the maximum 13. Fecundity differed between parishes (χ_4^2 =39.12,



Figure 1. Probability of surviving to adulthood (age 15) according to the sex of elder offspring (mean \pm s.e.). Values are adjusted means from the final model.



Figure 2. Lifetime reproductive success (number of children raised to age 15) according to the sex of elder offspring (mean \pm s.e.). Values are adjusted means from the final model.

p < 0.0001) and was higher for those in the rich and middle social classes than those in the poor class ($\chi_2^2 = 42.70$, p < 0.0001, adjusted means \pm s.e. 4.81 ± 0.18 , 4.66 ± 0.21 and 2.45 ± 0.28 for rich, middle class and poor, respectively). After controlling for these effects, we found that lifetime fecundity was significantly higher (12%) in those born after elder sisters than in those born after elder brothers



Figure 3. (a) Lifetime fecundity (number of children produced) according to sex of elder sibling (mean \pm s.e.). (b) Offspring survival rate (proportion of offspring surviving to age 15) according to the sex of elder offspring (mean \pm s.e.). Values are adjusted means from the final model.

 $(\chi_1^2=5.06, p=0.025;$ figure 3*a*). Lifetime fecundity was not significantly associated with year of birth $(\chi_1^2=0.11, p=0.74)$, birth order $(\chi_1^2=0.00, p=0.96)$, family size $(\chi_1^2=1.22, p=0.27)$, previous birth interval $(\chi_1^2=0.82, p=0.37)$ or own sex $(\chi_1^2=0.10, p=0.75)$.

After excluding childless individuals, 557 were included in the analysis of offspring survival rate. Only 25% of couples successfully raised all of the children they produced. Offspring survival rate, weighted in the analysis by the proportion of offspring successfully followed, differed between parishes (χ_4^2 =32.02, p<0.001) and was higher for those in the rich and middle social classes than those in the poor class. After controlling for these confounders, there was a non-significant trend towards offspring survival rate being higher for those born after elder female versus male siblings (χ_1^2 =2.80, p=0.094; figure 3b). Offspring survival rate was not associated with year of birth (χ_1^2 =2.68, p=0.10), birth order (χ_1^2 =1.97, p=0.16), family size (χ_1^2 =0.35, p=0.55), F₁ birth interval (χ_1^2 =2.17, p=0.14) or own sex (χ_1^2 =0.00, p=0.97).

4. DISCUSSION

Our results show that F_1 individuals who were born to mothers who had previously produced a son, when compared with those born to mothers who had previously produced a daughter, had similar probability of survival to adulthood but lower LRS. This latter result was mediated primarily through reduced lifetime fecundity of those born after an elder brother. Our results were apparent in all five parishes and across all social classes, indicating that they were not dependent on any particular ecological or social factors (see electronic supplementary material).

We consider three explanations to account for these observations. First, direct sibling interaction could lead to the reduced success of the younger offspring through the elder male siblings having a direct negative influence on the development of their younger siblings. As has been seen in previous studies on humans, elder offspring may influence the survival (Harpending & Pennington 1991), or reproductive success (Mace 1996) of their younger siblings. In particular, sex-specific competition between siblings for resources and reproductive opportunity may be an important determinant of reproductive success (Low 1990; Low & Clarke 1991; Mace 1996; Mulder 1998). Second, the results could be caused by nonindependence of the subsequent and previous offspring sex. Maternal condition may simultaneously influence both offspring LRS and the likelihood of producing male versus female offspring, specifically with mothers with greater access to resources being more likely to produce sons (Trivers & Willard 1973). Finally, individuals may have reduced success because their mother producing a son versus a daughter in the previous reproductive event rendered her less able to invest in the next offspring.

The hypothesis that males have a direct influence on their younger siblings predicts that if an elder sibling died before the birth of the focal individual, then there would be no relationship between elder sibling sex and the survival and LRS of the focal individual. Owing to high infant mortality, elder sibling death prior to the birth of the focal individual occurred in at least 30% of the cases in all datasets. It was therefore possible to test this hypothesis by entering into each model an interaction between elder sibling sex and whether the elder sibling died prior to or after the focal individual's birth. This interaction did not approach significance in either survival to age 15 ($\chi_1^2 =$ 1.54, p=0.21) or LRS ($\chi_1^2=0.32$, p=0.57). Similarly, the survival of the elder sibling to age 15 did not modify the relationships ($\chi_1^2 = 1.75$, p = 0.19; $\chi_1^2 = 0.47$, p = 0.49), pointing out that there was no indication of sex-specific post-natal interactions between siblings affecting our measures of offspring success. Finally, controlling for the completed total family size, which is an overall measure of sibling competition (Parker et al. 2002), failed to change the results. This suggests that the relationship of elder sibling sex with LRS and lifetime fecundity was not due to direct interaction between the elder sibling and the focal individual.

The hypothesis that non-random sex allocation might give rise to younger siblings born after sons having reduced success predicts that P offspring sex is influenced by maternal condition, with mothers who have access to a high level of resources being more likely to produce a son than a daughter (Trivers & Willard 1973). However, if maternal condition had influenced offspring sex, we would expect those born to mothers who had previously produced daughters to have lower than average LRS and lifetime fecundity. We would also expect LRS to be lowest in those cases where both the elder sibling and the focal individual were females and highest in those cases where both were males. This is because high-resource mothers would (i) benefit from a male bias in offspring sex allocation and (ii) produce high-quality offspring. However, not only was there no difference in LRS between the sexes but there was also no indication of any interactions between the sex of the focal individual and the elder sibling in these models (survival: $\chi_1^2=0.86$, p=0.35; LRS: $\chi_1^2=0.01$, p=0.93; see graphs in the electronic supplementary material). Therefore, nonindependence of offspring sex or sex ratio with respect to offspring LRS is unlikely to account for the strong relationship we observe between previous offspring sex and focal offspring LRS.

The simplest explanation for offspring born after sons having reduced reproductive success is that mothers who produced sons incurred a greater reproductive cost, and hence were less able to invest in future reproductive events, with this reduced investment having long-term consequences for her subsequent offspring. This conclusion is also supported by considerable evidence from other mammal studies, showing that producing sons versus daughters is indeed more costly to mothers as well as to subsequent offspring (table 1). Lower LRS and fecundity of those born after sons in this population can therefore be regarded as part of an emerging picture of a sex bias in the cost of offspring to mothers (see §1). For example, previous studies have shown that human offspring born after sons have reduced birth weight compared with those born after daughters (Trotnow et al. 1976; Magnus et al. 1985; Blanchard & Ellis 2001; Côté et al. 2003) and are also smaller as adults (I. J. Rickard 2007, unpublished results). Smaller sizes at birth and in adulthood are in turn related to reduced reproductive success in adulthood (Lummaa & Clutton-Brock 2002), offering one potential mechanism for our findings. As with all long-term correlative studies, special attention should be given to the potential of bias being introduced by the loss of individuals to follow up (Van de Pol & Verhulst 2005). Those individuals who were excluded from the analysis because they lacked specific data may have been affected differently by the sex of the offspring their mother produced previously. However, we consider this possibility unlikely, because these excluded individuals did not appear to differ from those included in terms of their discernible or life-history characteristics. Furthermore, the probability of exclusion from the data subsets was not related to elder sibling sex (comparison of proportions of those with elder male and female siblings between core data and subset 1, $\chi_1^2 = 0.58$, p = 0.45).

The consistency of the relationship between elder sibling sex and LRS is worth remarking upon. The association was not modified by mother's social class (two-level factor, $\chi_1^2=1.15$, p=0.29) or the interval separating the birth of the focal individual and their elder sibling ($\chi_1^2=0.78$, p=0.76). This is intriguing, because, for example, mothers of a high social class would have more access to resources, and therefore would be expected to withstand any higher energetic cost of producing a son over a daughter. Similarly, it would be expected that those mothers who gave birth to the focal individual a long time after the elder sibling would have been better able to recover from the extra cost of producing a son. However, in addition to, or perhaps instead of, their possibly higher demand on limited maternal resources, the cost differential in producing sons may be due to the factors that are independent of the mother's condition. For example, during development, male and female foetuses produce different quantities of hormones (Clark et al. 1991). Because these hormones may diffuse across foetal membranes and amniotic fluid, the endocrine profiles of expectant mothers may differ according to the sex of the offspring being carried. A hormone such as testosterone, which occurs in much higher concentration in a male foetus than in a female adult (Meulenberg & Hofman 1991), could interfere directly with a mother's ability to provide for future offspring by compromising her immunocompetence and increasing her susceptibility to costly disease (Klein 2000). This could in turn reduce her ability to invest in subsequent offspring. Another intriguing observation that has implications for evaluating the potential mechanism is that the effect is not conditional upon, or altered by, the sex of the focal individual. ($\chi_1^2 = 0.01$, p=0.93). It suggests that, while in polygynous ungulates the early determinants of reproductive success may differ between the sexes (Kruuk et al. 1999), in monogamous human societies such as this one, there may be less of a difference. To further understand the mechanism(s) behind our findings, we are currently investigating the relationship between elder sibling sex and those lifehistory traits that underlie lifetime fecundity.

Regardless of the underlying mechanism, we have shown clear evidence for an effect of previous maternal investment in a son on the LRS and lifetime fecundity of subsequent offspring. This has implications for calculations of maternal fitness and for considering the selective pressures and ecological constraints relevant to the evolution of adaptive sex-ratio adjustment. In polygynous mammals, a prospective mother with access to more resources than her competitors may benefit from producing a son rather than a daughter if she can better use those resources to improve the reproductive success of a son (Trivers & Willard 1973). Conversely, a prospective mother with access to a relatively low level of resources will benefit by producing a daughter, because any son she produces will ultimately lose out in competition for mates with the sons produced by high-resource females (Trivers & Willard 1973). While our results are not at odds with the qualitative predictions of the Trivers-Willard (1973) hypothesis, they suggest that the benefits of such adaptive sex-ratio variation may need to be weighed against additional costs which manifest themselves later, in the attenuated reproductive success of subsequent offspring.

In this study, we have shown, what is to our knowledge, the first evidence of a long-term intergenerational cost of reproduction in a long-lived species. Many studies have tested the effects of manipulated offspring brood size on offspring (Lessells 1986; Hare & Murie 1992; Mappes *et al.* 1995; Kunkele 2000). While failure to find intergenerational costs of enlarged broods may sometimes be due to those costs being expressed within the breeding individual, another reason may be that they are expressed in offspring reproductive performance. Owing to dispersal and practical constraints of field studies, animals from artificially manipulated broods or litters cannot be easily followed into their adult lives. However, the general findings of the small number of studies to have attempted such a long-term approach (Gustafsson & Sutherland 1988; Blondel *et al.* 1998; Koskela 1998; Naguib *et al.* 2006) reinforce the possibility that such unexplored costs of reproduction are widespread. The present study is the first to find a cost of reproduction manifested in reduced LRS of offspring. Observational studies of long-term pedigree data such as this one provide a useful means with which to approach key life-history questions that cannot conveniently be answered with many vertebrate study systems.

We thank Aino Siitonen, Kimmo Pokkinen and Timo Verho for collection of the demographic dataset. Thanks to Natural Environment Research Council (I.J.R.), the Royal Society of London (V.L. and A.F.R.) and the Academy of Finland (V.L.) for funding, and Rebecca Atkinson, Esa Koskela, Louis Longley, Isabelle Charmantier and two anonymous reviewers for their helpful comments.

REFERENCES

- Barker, D. J. P. 1998 Mothers, babies and health in later life. Edinburgh, UK: Churchill Livingstone.
- Bérubé, C. H., Festa-Bianchet, M. & Jorgenson, J. T. 1996 Reproductive costs of sons and daughters in Rocky Mountain bighorn sheep. *Behav. Ecol.* 7, 60–68. (doi:10. 1093/beheco/7.1.60)
- Birgersson, B. 1998 Male-biased maternal expenditure and associated costs in fallow deer. *Behav. Ecol. Sociobiol.* 43, 87–93. (doi:10.1007/s002650050470)
- Blanchard, R. & Ellis, L. 2001 Birth weight, sexual orientation and the sex of preceding siblings. *J. Biosoc. Sci.* 33, 451–467. (doi:10.1017/S0021932001004515)
- Blondel, J., Maistre, M., Perret, P., Hurtrez-Boussès, S. & Lambrects, M. M. 1998 Is the small clutch size of a Corsican blue tit population optimal? *Oecologica* 117, 80–89. (doi:10.1007/s004420050634)
- Clark, M. M., Crews, D. & Galef Jr, B. G. 1991 Concentrations of sex steroid hormones in pregnant and fetal Mongolian gerbils. *Physiol. Behav.* 49, 239–243. (doi:10.1016/0031-9384(91)90038-P)
- Clutton-Brock, T. H. 1991 *The evolution of parental care.* Monographs in behavior and ecology. Princeton, NJ: Princeton University Press.
- Clutton-Brock, T. H., Albon, S. D. & Guinness, F. E. 1981 Parental investment in male and female offspring in polygynous mammals. *Nature* **289**, 487–489. (doi:10. 1038/289487a0)
- Côté, K., Blanchard, R. & Lalumière, M. L. 2003 The influence of birth order on birth weight: does the sex of preceding siblings matter? *J. Biosoc. Sci.* 35, 455–462. (doi:10.1017/S0021932003004553)
- Drent, R. H. & Daan, S. 1980 The prudent parent: energetic adjustments in avian breeding. *Ardea* 68, 225–252.
- Festa-Bianchet, M. 1989 Individual differences, parasites, and the costs of reproduction for bighorn ewes (*Ovis* canadensis). J. Anim. Ecol. 58, 785–795. (doi:10.2307/5124)
- Festa-Bianchet, M., Jorgenson, J. T. & Réale, D. 2000 Early development, adult mass and reproductive success in bighorn sheep. *Behav. Ecol.* **11**, 633–639. (doi:10.1093/ beheco/11.6.633)
- Gomendio, M., Clutton-Brock, T. H., Albon, S. D., Guinness, F. E. & Simpson, M. J. 1990 Mammalian sex ratios and variation in costs of rearing sons and daughters. *Nature* 343, 261–263. (doi:10.1038/343261a0)
- Gustafsson, L. & Sutherland, W. J. 1988 The costs of reproduction in the collared flycatcher *Ficedula albicollis*. *Nature* 335, 813–815. (doi:10.1038/335813a0)

- Hare, J. F. & Murie, J. O. 1992 Manipulation of litter size reveals no cost of reproduction in Columbian groundsquirrels. *J. Mammal.* 73, 449–454. (doi:10.2307/ 1382083)
- Harpending, H. C. & Pennington, R. 1991 Age structure and sex-biased mortality among Herero pastoralists. *Hum. Biol.* 63, 329–353.
- Helle, S., Lummaa, V. & Jokela, J. 2002 Sons reduced maternal longevity in preindustrial populations. *Science* 296, 1085. (doi:10.1126/science.1070106)
- Hewison, A. J. M. & Galliard, J. M. 1999 Successful sons or advantaged daughters? The Trivers–Willard model and sexbiased maternal investment in ungulates. *Trends Ecol. Evol.* 14, 229–234. (doi:10.1016/S0169-5347(99)01592-X)
- Hogg, J. T., Hass, C. C. & Jenni, D. A. 1992 Sex-biased maternal expenditure in Rocky Mountain bighorn sheep. *Behav. Ecol. Sociobiol.* **31**, 243–251. (doi:10.1007/ BF00171679)
- Karn, M. & Penrose, L. S. 1952 Birth weight and gestation time in relation to maternal age, parity and infant survival. *Ann. Eugenics* 16, 147–164.
- Karskela, S. 2001 *Sukututkijan tietokirja*. Saarijärvi, Finland: Gummerus Kirjapaino Oy.
- Klein, J. L. 2000 The effects of hormones on sex differences in infection: from genes to behaviour. *Neurosci. Biobehav. Rev.* 24, 627–638. (doi:10.1016/S0149-7634(00)00027-0)
- Korpelainen, H. 2003 Human life histories and the demographic transition: a case study from Finland, 1870–1949. Am. J. Phys. Anthropol. 120, 384–390. (doi:10.1002/ajpa.10191)
- Koskela, E. 1998 Offspring growth, survival and reproductive success in the bank vole: a litter size manipulation experiment. *Oecologica* **115**, 379–384. (doi:10.1007/s004420050531)
- Kruuk, L. E. B., Clutton-Brock, T. H., Rose, K. E. & Guinness, F. E. 1999 Early determinants of lifetime reproductive success differ between the sexes in red deer. *Proc. R. Soc. B* 266, 1655. (doi:10.1098/rspb.1999.0828)
- Kunkele, J. 2000 Effects of litter size on the energetics of reproduction in a highly precocial rodent, the guinea pig. *J. Mammal.* 81, 691–700. (doi:10.1644/1545-1542(2000) 081<0691:EOLSOT>2.3.CO;2)
- Lack, D. 1947 The significance of clutch size. *Ibis* 89, 302–352.
- Lack, D. 1948a The significance of clutch size. Ibis 90, 25-45.
- Lack, D. 1948b The significance of litter size. *J. Anim. Ecol.* **17**, 45–50. (doi:10.2307/1608)
- Landete-Castillejos, T., García, A., López-Serrano, F. R. & Gallego, L. 2004 Maternal quality and differences in milk production and composition for male and female Iberian red deer calves (*Cervus elaphus hispanicus*). *Behav. Ecol. Sociobiol.* 57, 267–274. (doi:10.1007/s00265-004-0848-8)
- Lessells, C. M. 1986 Brood size in Canada geese: a manipulation experiment. *J. Anim. Ecol.* 55, 669–689. (doi:10.2307/4747)
- Low, B. S. 1990 Occupational status, landownership, and reproductive behavior in 19th-century Sweden: Tuna parish. Am. Anthropol. 92, 457–468. (doi:10.1525/aa. 1990.92.2.02a00130)
- Low, B. S. & Clarke, A. L. 1991 Family patterns in nineteenth-century Sweden. J. Family Hist. 16, 117–138.
- Lummaa, V. & Clutton-Brock, T. H. 2002 Early development, survival and reproduction in humans. *Trends Ecol. Evol.* 17, 141–147. (doi:10.1016/S0169-5347(01)02414-4)
- Lummaa, V., Pettay, J. E. & Russell, A. F. 2007 Male twins reduce fitness of female co-twins in humans. *Proc. Natl Acad. Sci. USA* **104**, 10 915–10 920. (doi:10.1073/pnas. 0605875104)

- Mace, R. 1996 Biased parental investment and reproductive success in Gabbra pastoralists. *Behav. Ecol. Sociobiol.* 38, 75–81. (doi:10.1007/s002650050219)
- Mace, R. & Sear, R. 1997 Birth interval and the sex of children in a traditional African population: an evolutionary analysis. *J. Biosoc. Sci.* 29, 499–507. (doi:10.1017/ S0021932097004999)
- Magnus, P., Berg, K. & Bjerkedal, T. 1985 The association of parity and birth weight: testing the sensitization hypothesis. *Early Hum. Dev.* 12, 49–54. (doi:10.1016/ 0378-3782(85)90136-7)
- Mappes, T., Koskela, E. & Ylonen, H. 1995 Reproductive costs and litter size in the bank vole. *Proc. R. Soc. B* 261, 19–24. (doi:10.1098/rspb.1995.0111)
- Meulenberg, P. M. M. & Hofman, J. A. 1991 Maternal testosterone and fetal sex. *J. Steroid Biochem. Mol. Biol.* 39, 51–54. (doi:10.1016/0960-0760(91)90012-T)
- Mulder, M. B. 1998 Brothers and sisters—how sibling interactions affect optimal parental allocations. *Hum. Nat.* 9, 119–161. (doi:10.1007/s12110-998-1001-6)
- Naguib, M., Nemitz, A. & Gil, D. 2006 Maternal developmental stress reduces reproductive success of female offspring in zebra finches. *Proc. R. Soc. B* 273, 1901–1905. (doi:10.1098/rspb.2006.3526)
- Parker, G. A., Royle, N. J. & Hartley, I. R. 2002 Intrafamilial conflict and parental investment: a synthesis. *Phil. Trans. R. Soc. B* 357, 295–307. (doi:10.1098/rstb. 2001.0950)
- Pettay, J. E., Helle, S., Jokela, J. & Lummaa, V. 2007 Natural selection on female life-history traits in relation to socioeconomic class in pre-industrial human populations. *PLoS ONE* 2, e606. (doi:10.1371/journal.pone.0000606)

- Phillips, D. I. W., Handelsman, D. J., Eriksson, J. G., Forsén, T., Osmond, C. & Barker, D. J. P. 2001 Prenatal growth and subsequent marital status: longitudinal study. *Br. Med. J.* 322, 771. (doi:10.1136/bmj.322.7289.771)
- Roff, D. A. 1992 *The evolution of life histories: theory and analysis.* New York, NY: Chapman and Hall.
- SAS 1990 SAS/STAT user's guide. Cary, NC: SAS Institute, Inc.
- Tamimi, R., Lagiou, P., Mucci, L. A., Hsieh, C., Adami, H. & Trichopoulos, D. 2003 Average energy intake among pregnant women carrying a boy compared with a girl. *Br. Med. J.* 326, 1245–1547. (doi:10.1136/bmj.326.7401. 1245)
- Trivers, R. L. & Willard, D. E. 1973 Natural selection of parental ability to vary the sex ratio of offspring. *Science* **179**, 90–92. (doi:10.1126/science.179.4068.90)
- Trotnow, S., Bregulla, K. & Flügel, K. 1976 Untersuchung über die Körpergröβe und das Körpergewicht von Neugeborenen in Abhängigkeit vom Paritätsstatus [Research on the body size and body weight of newborns as a function of maternal parity]. *Geburtshilfe Frauenheilkunde* **36**, 744–750.
- Vågerö, D. & Modin, B. 2002 Prenatal growth, subsequent marital status, and mortality: longitudinal study. *Br. Med. J.* 324, 398. (doi:10.1136/bmj.324.7334.398)
- Van de Pol, M. & Verhulst, S. 2005 Age-dependent traits: a new statistical model to separate within- and betweenindividual effects. Am. Nat. 167, 766–773. (doi:10.1086/ 503331)
- van Noordwijk, A. J. & Dejong, G. 1986 Acquisition and allocation of resources—their influence on variation in lifehistory tactics. *Am. Nat.* **128**, 137–142. (doi:10.1086/ 284547)