

Effects of early developmental conditions on innate immunity are only evident under favourable adult conditions in zebra finches

Greet De Coster · Simon Verhulst · Egbert Koetsier ·
Liesbeth De Neve · Michael Briga · Luc Lens

Received: 15 July 2011 / Revised: 27 October 2011 / Accepted: 30 October 2011 / Published online: 12 November 2011
© Springer-Verlag 2011

Abstract Long-term effects of unfavourable conditions during development can be expected to depend on the quality of the environment experienced by the same individuals during adulthood. Yet, in the majority of studies, long-term effects of early developmental conditions have been assessed under favourable adult conditions only. The immune system might be particularly vulnerable to early environmental conditions as its development, maintenance and use are thought to be energetically costly. Here, we studied the interactive effects of favourable and unfavourable conditions during nestling and adult stages on innate immunity (lysis and agglutination scores) of captive male and female zebra finches (*Taeniopygia guttata*). Nestling environmental conditions were manipulated by a brood size experiment, while a foraging cost treatment was imposed on the same individuals during adulthood. This combined treatment showed that innate immunity of adult zebra finches is affected by their early developmental conditions and varies between both sexes. Lysis scores, but not agglutination scores, were higher in individuals raised in small broods and in males. However, these effects were only present in birds that experienced low foraging costs. This study shows that the quality of the

adult environment may shape the long-term consequences of early developmental conditions on innate immunity, as long-term effects of nestling environment were only evident under favourable adult conditions.

Keywords Brood size manipulation · Complement · Developmental stress · Foraging costs manipulation · Immunocompetence · Natural antibodies

Introduction

Adverse conditions experienced during early life, such as suboptimal nest temperature, nutritional stress and exposure to pathogens and toxins, have been shown to negatively affect an individual's physiology, survival and reproductive success during adulthood in a wide range of species (e.g. Gluckman and Hanson 2004; Lindström 1999; Lummaa and Clutton-Brock 2002; Metcalfe and Monaghan 2001). Long-term consequences of early developmental conditions have, however, been found to vary with the quality of the adult environment in opposing ways (reviewed in Monaghan 2008). First, it has been demonstrated that nestlings raised under poor conditions suffer a suboptimal development and experience negative effects irrespective of the quality of the adult environment (Grafen 1988; Monaghan 2008). Second, several studies found that long-term effects of developmental stress remain masked under favourable adult conditions as physiological responses are not affected by an individual's quality (e.g. Hendrickx et al. 2003; Lens et al. 2002; Talloen et al. 2010). However, when environmental conditions deteriorate and individuals become challenged, qualitative differences become reflected in their physiological responses, ultimately leading to variation in survival (e.g. Hendrickx et al. 2003; Lens et al. 2002; Talloen et al. 2010). Third,

Communicated by: Sven Thatje

G. De Coster (✉) · L. De Neve · L. Lens
Department of Biology, Terrestrial Ecology Unit,
Ghent University,
K.L. Ledeganckstraat 35,
9000 Ghent, Belgium
e-mail: greet.decoester@ugent.be

S. Verhulst · E. Koetsier · M. Briga
Behavioural Biology, Centre for Life Sciences,
University of Groningen,
PO Box 11103, 9700 CC Groningen, The Netherlands

individuals developing under unfavourable circumstances may cope better with poor conditions during adulthood because of the similarity between early and later environments. If this is the case, a reverse pattern can be expected when adult conditions are more favourable, as organisms raised in poor environments are not developmentally adapted to more favourable ambient conditions (Monaghan 2008).

As the development, maintenance and use of an organism's immune system are assumed to be energetically costly (e.g. Klasing 2004; Lochmiller and Deerenberg 2000; Norris and Evans 2000; Sheldon and Verhulst 1996), immune function can be expected to be suppressed in large broods because of higher nutritional stress and increased sibling competition (e.g. Naguib et al. 2004; Saino et al. 1997). Yet, poor immunity can come at a price as the immune system is the main physiological mechanism to resist disease, and hence, to safeguard future survival and reproduction (reviewed in Møller and Saino 2004). This might be particularly true for two components of the constitutive innate immune system, the natural antibodies (hereafter NABs) and complement cascade, which provide the first defence mechanisms against a broad spectrum of pathogens, and therefore seem to play a crucial role in enhancing an organism's survival (Ochsenbein and Zinkernagel 2000). NABs (mostly IgM) are already present in individuals that have not been exposed to antigens (Bandeira et al. 1988). Their function is to bind to a particular antigen or pathogen, to clear foreign substances and to initiate the complement system, which consists of circulating serum proteins that are sequentially activated to result in cell lysis (Boes 2000 and references therein). Although NABs and the complement cascade are assumed to interact, their concentrations are usually not, or only weakly, correlated (De Coster et al. 2010; Mendes et al. 2006; Møller and Haussy 2007; Soler et al. 2007, but see Parejo et al. 2007). Despite the theoretical expectations outlined above, empirical evidence of the effects of early developmental conditions on innate immune function remains scarce and heterogeneous. For example, while complement activation was positively related to clutch size in a suite of tropical bird species, no such effect was found for NABs (Lee et al. 2008). In a study on European rollers (*Coracias garrulus*), both components varied with brood hierarchy in nestling body mass, though in opposite ways. While lighter nestlings had more NABs compared to heavier ones, the former showed less complement activity (Parejo et al. 2007). Given the role of constitutive innate immunity as a first-line defence against infections and its effects on survival (Ochsenbein and Zinkernagel 2000), it is important to gain more general insights into the long-term effects of developmental conditions, in addition to their interaction with the adult environment, on this branch of the immune system.

Furthermore, it remains unexplored whether such effects on innate immunity differ between the sexes, despite indications that inflammatory immune responses in reaction to conditions during development and adulthood differ between male and female mallard ducks (*Anas platyrhynchos*; Butler and McGraw 2011).

Here, we attempt to explore this research question by experimentally manipulating environmental conditions during the nestling and adult stages in captive male and female zebra finches (*Taeniopygia guttata* Vieillot) and study effects on their innate immune capacity. Brood size manipulations are a well-known technique to manipulate nutritional intake and the level of competition of individual nestlings (Griffith and Buchanan 2010). At a short term, being raised in a large brood reduces body sizes, growth rates, phytohaemagglutinin (PHA) inflammatory responses and survival of nestlings (e.g. Alonso-Alvarez et al. 2006; de Kogel 1997b; Naguib et al. 2004; Tschirren et al. 2009), while at a longer term, it affects adult body size, metabolic rates, inflammatory responses and reproductive success (e.g. Alonso-Alvarez et al. 2006; Tschirren et al. 2009; Verhulst et al. 2006). Early developmental conditions can further be expected to differentially affect adult innate immunity in male and female zebra finches, as the latter have been shown to be more negatively affected by harsh rearing conditions (e.g. Bradbury and Blakey 1998; de Kogel 1997b; Kilner 1998; Martins 2004; Verhulst et al. 2006), and inflammatory and humoral immune responses have been shown to vary between sexes and contexts (Love et al. 2008; McGraw and Ardia 2005). Furthermore, increased foraging costs in adulthood result in physiological adjustments to save energy (Wiersma et al. 2005; Wiersma and Verhulst 2005) and might therefore also affect innate immunity because of the associated energetic costs (Klasing 2004).

The majority of studies on long-term fitness effects of development under poor conditions were conducted under favourable adult conditions (reviewed in Monaghan 2008), which precludes the evaluation of possible interactions between early and later life conditions. Here, we consider favourable and unfavourable adult conditions and study the interactive effects with manipulations of early condition. Nestling environmental conditions were manipulated through a brood size experiment, while the same individuals were subsequently subjected to a foraging cost treatment during adulthood. Based on these paired experiments, we examined if, and to what extent: (a) environmental conditions during the nestling and/or adult stage have persistent effects on adult innate immune responses; (b) long-term effects of nestling environment depend on the quality of the adult environment and (c) effects of environmental conditions differ between males and females.

Material and methods

Bird husbandry and experimental design

Our study was conducted between 2006 and 2009 in a captive population kept at the University of Groningen, The Netherlands. All individuals were hatched under standard laboratory conditions and subjected to a cross-fostering at the age of 5 days such that nestlings were either raised in a small brood (two nestlings) or a large brood (six nestlings). In this procedure, we ensured that parents rearing small or large broods did not on average differ in reproductive traits such as clutch size or brood size before manipulation. We can therefore safely assume that parents rearing small or large broods did not on average differ in their ability to care for a brood. Birds from which the brood size had not been manipulated (i.e. comprising two or six nestlings in the original brood) were excluded from our analyses. Experimental brood sizes fell within the range normally observed in wild (Zann 1996) and laboratory (Naguib et al. 2004) populations. We manipulated the brood size towards the extremes of the normal range in order to trigger sufficient variation in nestling environment between the two experimental groups. When individuals were between 3 and 23 months old, they were distributed over eight single-sex outdoor aviaries (four with males only, four with females only). Birds were kept in same sex groups because the opportunity to invest in sexual interactions could mitigate treatment effects on physiology and life span when birds in the poor foraging environment (see further) engage less in such interactions. Individuals were divided equally over the aviaries with respect to the experimental brood sizes and their age. In this way, each aviary contained 32–37 individuals of different age and brood size treatments and was exposed to a natural light–dark cycle and ambient temperature. While drinking water and tropical food seed mixture was supplied ad libitum in each aviary, foraging costs per reward were manipulated by placing perches under the food container in four cages (two per sex; *low foraging cost* treatment) while not placing them in four other cages (two per sex; *high foraging cost* treatment). This treatment forced birds to fly back and forth between a perch and the food container for each seed they consumed. The seeds that fell from the seed containers were not accessible to prevent individuals from each aviary to forage on spilled seed at low cost. Previous work on the same birds showed that birds experiencing high foraging costs spent more than twice as much time foraging and significantly reduced basal metabolic rates compared to low foraging cost birds, demonstrating the effectiveness of the imposed treatment to increase foraging costs and to affect physiological responses (Koetsier and Verhulst 2011). Upon blood collection, survival rates were lower in

high-cost foragers raised in large broods (39 out of 69 individuals, 57%) compared to those raised in small broods (53 out of 64, 83%) ($P=0.003$). Rates did not differ among low-cost foragers (small, 42 out of 58, 72%; large, 50 out of 75, 67%; $P=0.50$), while the interaction between brood size and foraging costs was marginally significant ($P=0.077$; generalised linear mixed model; see below for random effects). A total of 96 females and 88 males that were still alive in October 2009 (16–45 months old and 8–22 months subjected to the foraging cost treatment) were weighed, and 100–150 μ l blood was collected in heparinized capillary tubes via brachial vein puncture with a sterile 25-gauge hypodermic needle. Individuals raised in large broods had a lower body mass during adulthood than individuals raised in small broods ($P=0.05$; linear mixed model; see below for details), while individuals subjected to the high foraging cost treatment had a lower body mass than individuals subjected to the low-cost treatment ($P=0.01$). These results validate our brood size and foraging cost treatment.

Haemolysis–haemagglutination assay

Blood samples were immediately transferred to a microfuge tube and kept cool before being centrifuged (5 min at $10,000\times g$). The plasma was separated from the cells and frozen at -20°C . The levels of circulating NABs and complement were estimated by means of the haemolysis–haemagglutination assay developed by Matson et al. (2005) within 10 days after blood collection. This assay is based on red blood cell agglutination and NAB-mediated complement activation. Quantification is done by a serial dilution (1:2) of plasma samples, starting with 50 μ l plasma, with phosphate-buffered saline in a 96-well assay plate, which was then incubated with the same amount of rabbit red blood cell suspension for 90 min at 37°C . Titres were scored blindly with respect to experimental brood size, foraging cost treatment, sex and age as the highest dilution step at which agglutination (NAB) and lysis (complement) of red blood cells were observed. Higher scores were associated with stronger immune responses. Half scores were assigned when the termination of the reaction was intermediate between two wells.

Statistical analysis

Agglutination scores ranged between 3 and 8 (mean \pm SD= 4.65 ± 0.73) and were treated as samples from a normal distribution. Lysis scores ranged between 0 and 2 (mean \pm SD= 0.17 ± 0.36) and were treated as samples from a Poisson distribution to allow for the non-negative values for lysis scores despite the small mean. While lysis scores cannot be treated as count data *sensu stricto*, noninteger response variables can be included when modelling

Poisson-distributed data in SAS 9.2 (SAS Institute, 2002–2008; see also Dugdale et al. 2010). Linear mixed models with restricted maximum likelihood and Kenward–Roger approximation for the degrees of freedom (Kenward and Roger 1997) were used to model agglutination scores. Generalised linear mixed models with logarithmic link function were fitted to model lysis scores. Starting models contained fixed factors brood size (small or large), foraging costs (high or low) and sex, all two-factor interactions and the three-factor interaction. Models were corrected for the covariates age, body mass, treatment duration and original brood size (i.e. in the nest of origin) to account for differences in age, body mass, time involved in the foraging cost treatment and brood size shift, respectively. Body condition index (calculated as the residual of the linear regression of body mass on tarsus length with both measures being logarithmic transformed) was not included in the models to avoid multi-collinearity problems due to its high correlation with body mass ($\rho=0.93$; $P<0.0001$). Nest of rearing and aviary were included as random factors in all models to avoid pseudoreplication from sharing a common nest and adult environment. Non-significant fixed effects were removed from the model following a backward stepwise procedure. All analyses were performed with procedures Mixed (agglutination) and Glimmix (lysis) in SAS 9.2.

Results

Overall, lysis scores did not differ between foraging cost treatments, brood sizes or sexes (all $P>0.19$). However, the lysis scores significantly differed between brood sizes and sexes when accounting for the foraging cost treatment that adults were exposed to (interactions between brood size and foraging costs $P=0.021$, and between sex and foraging costs $P=0.022$; Table 1). Under high foraging costs, the lysis scores did not differ between individuals raised in large and small broods (post hoc test, $t_{54}=-1.21$; $P=0.23$; Fig. 1a) or between males and females (post hoc test, $t_{54}=0.96$; $P=0.34$; Fig. 1b). In contrast, among low-cost foragers, lysis scores were significantly smaller in individuals raised in large broods (post hoc test, $t_{54}=2.12$; $P=0.038$; Fig. 1a) and in females (post hoc test, $t_{54}=-2.29$; $P=0.026$; Fig. 1b).

In accordance with the lysis scores, average agglutination scores did not vary with the brood size manipulation and foraging cost treatment, nor did they differ between the sexes (all $P>0.19$). However, in contrast to the former, the effect of the foraging treatment did not depend on whether individuals had been raised in small or large broods ($P=0.71$). Likewise, there was no effect of either treatment (or their interaction) when the

Table 1 Generalised linear mixed model analysis of lysis scores in relation to brood size manipulation, foraging cost treatment and sex

Model term	df	F	P
Sex	1, 54	1.18	0.28
Brood size	1, 54	0.50	0.48
Foraging costs	1, 54	0.12	0.73
Sex \times brood size	1, 53	0.02	0.88
Sex \times foraging costs	1, 54	5.54	0.022
Brood size \times foraging costs	1, 54	5.64	0.021
Sex \times brood size \times foraging costs	1, 52	0.08	0.77

Significant terms are indicated in bold

df degrees of freedom

sex of the individuals was taken into account (all $P>0.57$). Lysis and agglutination scores were not related with age or body mass, neither did they vary with the time that an individual was involved in the foraging cost treatment or the original brood size (all $P>0.22$).

Discussion

By experimentally manipulating environmental conditions in nestlings (brood size experiment) and adults (foraging




Fig. 1 Effects of foraging cost manipulations on lysis scores in relation to (a) experimental brood size and (b) sex. Small (S) and large (L) broods and female (F) and male (M) birds are represented by filled and open bars, respectively. Numbers represent samples sizes; error bars represent standard errors

cost treatment), we showed that lysis titres, but not agglutination titres, were affected by nestling environmental conditions and sex, depending on the adult environment. Lysis scores of low-cost foragers (i.e. favourable adult condition) were significantly lower if these individuals had been raised in large broods, whereas the scores of high-cost foragers (i.e. unfavourable adult condition) were independent of brood size. Long-term negative effects of development under unfavourable conditions may either be expected (a) to be evident under both favourable and unfavourable adult conditions; (b) to be evident under unfavourable adult conditions only or (c) to be stronger under favourable adult conditions because of the mismatch between nestling and adult conditions (see [Introduction](#)). Though higher lysis scores under matching nestling and adult conditions (Fig. 1a) seem to provide support for the latter prediction, the non-significant effect of brood size under high foraging costs rather implies another mechanism, i.e. unfavourable nestling conditions causing long-term negative effects on (at least one aspect of) an individual's physiology but being evident under favourable adult conditions only. It has earlier been shown that increased foraging costs result in physiological adjustments and decreased investment in somatic maintenance to save energy (Wiersma et al. 2005; Wiersma and Verhulst 2005). Therefore, birds experiencing high foraging costs might simply have withdrawn energy from the maintenance of innate immunity because of the associated energetic costs (Klasing 2004). If so, long-term effects of early developmental conditions might be masked because of the reallocation of the energy budget.

Alternatively, it is also possible that effects under unfavourable conditions were masked due to differential mortality. High-cost foragers (Wiersma et al. 2005; Wiersma and Verhulst 2005) and individuals raised in large broods (e.g. Alonso-Alvarez et al. 2006; Tschirren et al. 2009; Verhulst et al. 2006) usually experience negative effects on a variety of fitness-related traits, in particular under unfavourable adult conditions (e.g. Hendrickx et al. 2003; Lens et al. 2002; Talloen et al. 2010). Indeed, high-cost foragers suffered from higher mortality when raised in large broods, whereas a similar effect was not apparent in low-cost foragers. If differential mortality under a combined 'large brood/high foraging cost' treatment was biased towards individuals with a low immune response (Møller and Saino 2004), the average immune response of surviving (and thus blood sampled) adults would be expected to increase and long-term effects on immunity of high-cost foragers to be masked. Unfortunately, we did not measure immune response when birds fledged in order to test this hypothesis. The fact that lysis scores in low-cost foragers (no bias in mortality) were smaller when raised in large broods supports a long-term cost of brood size on innate immunity. Our results contrast those of a study on free-

living great tits (*Parus major*) where (morphological) differences between individuals that developed in favourable and adverse conditions were only apparent under suboptimal conditions (Talloen et al. 2010). Yet, we propose that the same mechanism may be acting in both studies (i.e. adverse consequences of developmental stress being more severe in unfavourable adult conditions), but causing different effects in adulthood depending on whether or not differential mortality was involved.

Likewise, differential mortality between sexes may explain why female low-cost foragers had lower immune responses than males, while this sexual dimorphism in immune function was not evident when foraging entailed higher energetic costs. Given the higher mortality rates in female zebra finches under harsh environmental conditions (Bradbury and Blakey 1998; de Kogel 1997b; Kilner 1998), we suggest that mortality rates were biased towards females with relatively weak immune capacity such that the remaining females mounted higher immune responses, on average (see also Møller and Saino 2004). Alternatively, seeing that immunity is a highly plastic trait that varies sex-specifically and in a context-dependent way (Love et al. 2008), resource allocation trade-offs between immunity and other functions may differ between sexes. Male high-cost foragers may prioritise the investment in structures associated with mate acquisition, such as plumage ornaments, at the expense of immunocompetence to have similar mating success than individuals living in favourable conditions (Arnold et al. 2007; Naguib and Nemitz 2007). In contrast, females may rather maximise their survival through investments in immunity (Møller and Saino 2004). As a result, the immune responses of both sexes might be similar under poor environmental conditions.

The inverse effects of brood size on lysis titres of low-cost foragers are in agreement with the negative relationships of brood size with PHA response (Alonso-Alvarez et al. 2006; Naguib et al. 2004) earlier observed in zebra finch nestlings (but not in adults, see Tschirren et al. 2009). Several hypotheses have been invoked to explain these patterns. First, immune responses in large broods might be reduced as a result of the restricted amount of resources that nestlings receive from either parent (e.g. Fargallo et al. 2002; Hōrak et al. 1999; Saino et al. 1997). Second, lower immune responses in large broods might be caused by increased sibling competition resulting in an increase in testosterone (Naguib et al. 2004) and a suppressed immune function (Boonekamp et al. 2008; Fargallo et al. 2007; Folstad and Karter 1992; but see Roberts et al. 2004). Evidence for immunosuppressive effects of testosterone, however, is still lacking for zebra finches. Third, it has been suggested that stress caused by higher sibling competition increases corticosterone levels which, in turn, may suppress immunity (Saino et al. 2003). Correspondingly, prolonged

fasting resulted in increased corticosterone and decreased NAb levels in female mallards (Bourgeon et al. 2010). Yet, administering corticosterone did not alter innate immune capacity in barn owl (*Tyto alba*) nestlings (Stier et al. 2009), while brood size manipulation produced no effect on corticosterone in nestlings of the spotless starling (*Sturnus unicolor*; Gil et al. 2008), and have not been measured in zebra finches. However, as we only measured two components of the innate immune system, we cannot rule out that other branches of (innate) immunity were differently affected. It is possible that individuals optimised their investment in each component of the immune system, depending on the costs and benefits of investment in a particular component and on the conditions they encounter (Norris and Evans 2000).

Yet, the results of this study do not allow to fully exclude an alternative mechanism, i.e. that individuals raised in large broods possessed a lower ability to obtain food during adulthood. If so, the lower innate immune responses in individuals raised in large broods would not be caused by a long-term effect of nutritional conditions during early development but rather by a reduced resource consumption during adulthood and, as a result, a lower investment in the maintenance of the immune system. Yet, we think that such mechanism is unlikely to operate in our study population, as the time spent eating and dominance over food during adulthood are not affected by brood size in zebra finches (de Kogel 1997a) and body mass and lysis score were not correlated in this study. Furthermore, without knowledge of an individual's degree of pathogen infection, high immune responses can be interpreted in at least two different ways, i.e. as an indication of a well-functioning immune system (indicative of good individual condition) or as a consequence of intense pathogen infection (indicative of poor condition; Norris and Evans 2000). In the latter case, high lysis scores in birds from small broods would indicate that these birds are exposed to a higher number of infections. Yet, while NAb levels have been shown to rise in response to an experimentally increased level of parasite exposure, a similar effect was not found for complement activation (De Coster et al. 2010).

Unlike lysis scores, the agglutination scores were not affected by brood size or foraging cost treatments, nor associated with a bird's sex. Likewise, NAb concentrations did not vary with clutch size or food limitation in several bird species (e.g. Buehler et al. 2009a; Lee et al. 2008), but significant relationships have been found with growth rate, mass differences among nestlings, prolonged fasting, body mass and environmental conditions in other species (e.g. Bourgeon et al. 2010; Buehler et al. 2009b; Mauck et al. 2005; Parejo et al. 2007). The fact that agglutination titres did not vary with the experimental treatments in our study suggests that NAb levels are not (or only slightly) affected by

environmental conditions in zebra finches or that the species actively buffers NAb levels against stress-induced changes (see also Buehler et al. 2009a), possibly at the expense of investment in other life history traits. Further experimental study is needed to discriminate between both hypotheses.

To summarise, long-term effects of developmental conditions on physiology were only apparent under favourable conditions in adulthood. We showed that one component of innate immune function of adult zebra finches is affected by the conditions experienced during development and varies between sexes. To better understand the putative adaptive mechanisms underlying these relationships, future experiments that take into account resource allocation trade-offs among multiple condition-dependent traits are required.

Acknowledgements We thank M. Kouwen, F. Tamminga and M. Salomons for assistance in building and developing the design for the foraging costs experiment, E. Mulder for help with the haemolysis–haemagglutination assay and W. Müller and three anonymous reviewers for valuable comments on an earlier version of the manuscript. The experiments were performed in accordance to the ethical guidelines of the Animal Experimentation Committee of the University of Groningen (licence number 5150). G.D.C. was supported by a doctoral grant and by FWO research community WO.037.10N from the Research Foundation Flanders (FWO). E.K. and S.V. were supported by a Vici-grant to S.V. from The Netherlands Organisation for Scientific Research (NWO).

References

- Alonso-Alvarez C, Bertrand S, Devevey G, Prost J, Favre B, Chastel O, Sorci G (2006) An experimental manipulation of life-history trajectories and resistance to oxidative stress. *Evolution* 60:1913–1924. doi:10.1111/j.0014-3820.2006.tb00534.x
- Arnold KE, Blount JD, Metcalfe NB, Orr KJ, Adam A, Houston D, Monaghan P (2007) Sex-specific differences in compensation for poor neonatal nutrition in the zebra finch *Taeniopygia guttata*. *J Avian Biol* 38:356–366. doi:10.1111/j.2007.0908-8857.03818.x
- Bandeira A, Coutinho A, Martinez C, Pereira P (1988) The origin of “natural antibodies” and the internal activity in the immune system. *Int Rev Immunol* 3:47–58. doi:doi:10.3109/08830188809051181
- Boes M (2000) Role of natural and immune IgM antibodies in immune responses. *Mol Immunol* 37:1141–1149
- Boonekamp JJ, Ros AHF, Verhulst S (2008) Immune activation suppresses plasma testosterone level: a meta-analysis. *Biol Lett* 4:741–744. doi:10.1098/rsbl.2008.0347
- Bourgeon S, Kauffmann M, Geiger S, Raclot T, Robin J-P (2010) Relationships between metabolic status, corticosterone secretion and maintenance of innate and adaptive humoral immunities in fasted re-fed mallards. *J Exp Biol* 213:3810–3818. doi:10.1242/jeb.045484
- Bradbury RR, Blakey JK (1998) Diet, maternal condition, and offspring sex ratio the zebra finch, *Poephila guttata*. *Proc R Soc Lond B Biol Sci* 265:895–899. doi:10.1098/rspb.1998.0375
- Buehler DM, Encinas-Viso F, Petit M, Vezina F, Tieleman BI, Piersma T (2009a) Limited access to food and physiological trade-offs in

- a long-distance migrant shorebird. II. Constitutive immune function and the acute-phase response. *Physiol Biochem Zool* 82:561–571. doi:10.1086/603635
- Buehler DM, Tieleman BI, Piersma T (2009b) Age and environment affect constitutive immune function in red knots (*Calidris canutus*). *J Ornithol* 150:815–825. doi:10.1007/s10336-009-0402-6
- Butler MW, McGraw KJ (2011) Past or present? Relative contributions of developmental and adult conditions to adult immune function and coloration in mallard ducks (*Anas platyrhynchos*). *J Comp Physiol B* 181:551–563. doi:10.1007/s00360-010-0529-z
- De Coster G, De Neve L, Martín-Gálvez D, Thery L, Lens L (2010) Variation in innate immunity in relation to ectoparasite load, age and season: a field experiment in great tits (*Parus major*). *J Exp Biol* 213:3012–3018. doi:10.1242/jeb.042721
- de Kogel CH (1997a) Does brood size manipulation affect later competitive behaviour of offspring? *Behaviour* 134:161–172
- de Kogel CH (1997b) Long-term effects of brood size manipulation on morphological development and sex-specific mortality of offspring. *J Anim Ecol* 66:167–178
- Dugdale HL, Nouvellet P, Pope LC, Burke T, Macdonald DW (2010) Fitness measures in selection analyses: sensitivity to the overall number of offspring produced in a lifetime. *J Evol Biol* 23:282–292. doi:10.1111/j.1420-9101.2009.01896.x
- Fargallo JA, Laaksonen T, Poyri V, Korpimäki E (2002) Inter-sexual differences in the immune response of Eurasian kestrel nestlings under food shortage. *Ecol Lett* 5:95–101. doi:10.1046/j.1461-0248.2002.00290.x
- Fargallo JA, Martínez-Padilla J, Toledano-Díaz A, Santiago-Moreno J, Davila JA (2007) Sex and testosterone effects on growth, immunity and melanin coloration of nestling Eurasian kestrels. *J Anim Ecol* 76:201–209. doi:10.1111/j.1365-2656.2006.01193.x
- Folstad I, Karter AJ (1992) Parasites, bright males, and the immunocompetence handicap. *Am Nat* 139:603–622
- Gil D, Bulmer E, Celis P, Puerta M (2008) Increased sibling competition does not increase testosterone or corticosterone levels in nestlings of the spotless starling (*Sturnus unicolor*). *Horm Behav* 54:238–243. doi:10.1016/j.yhbeh.2007.11.013
- Gluckman PD, Hanson MA (2004) Living with the past: evolution, development, and patterns of disease. *Science* 305:1733–1736. doi:10.1126/science.1095292
- Grafen A (1988) On the uses of data on lifetime reproductive success. In: Clutton-Brock T (ed) *Reproductive success*. University of Chicago Press, Chicago, pp 454–471
- Griffith SC, Buchanan KL (2010) Maternal effects in the zebra finch: a model mother reviewed. *Emu* 110:251–267. doi:10.1071/mu10006
- Hendrickx F, Maelfait JP, Lens L (2003) Relationship between fluctuating asymmetry and fitness within and between stressed and unstressed populations of the wolf spider *Pirata piraticus*. *J Evol Biol* 16:1270–1279. doi:10.1046/j.1420-9101.2003.00633.x
- Hörak P, Tegelmann L, Ots I, Møller AP (1999) Immune function and survival of great tit nestlings in relation to growth conditions. *Oecologia* 121:316–322
- Kenward MG, Roger JH (1997) Small sample inference for fixed effects from restricted maximum likelihood. *Biometrics* 53:983–997
- Kilner R (1998) Primary and secondary sex ratio manipulation by zebra finches. *Anim Behav* 56:155–164. doi:10.1006/anbe.1998.0775
- Klasing KC (2004) The costs of immunity. *Acta Zool Sin* 50:961–969
- Koetsier E, Verhulst S (2011) A simple technique to manipulate foraging costs in seed-eating birds. *J Exp Biol* 214:1225–1229. doi:10.1242/jeb.050336
- Lee KA, Wikelski M, Robinson WD, Robinson TR, Klasing KC (2008) Constitutive immune defences correlate with life-history variables in tropical birds. *J Anim Ecol* 77:356–363. doi:10.1111/j.1365-2656.2007.01347.x
- Lens L, Van Dongen S, Matthysen E (2002) Fluctuating asymmetry as an early warning system in the critically endangered Taita thrush. *Conserv Biol* 16:479–487. doi:10.1046/j.1523-1739.2002.00516.x
- Lindström J (1999) Early development and fitness in birds and mammals. *Trends Ecol Evol* 14:343–348. doi:10.1016/S0169-5347(99)01639-0
- Lochmiller RL, Deerenberg C (2000) Trade-offs in evolutionary immunology: just what is the cost of immunity? *Oikos* 88:87–98. doi:10.1034/j.1600-0706.2000.880110.x
- Love OP, Salvante KG, Dale J, Williams TD (2008) Sex-specific variability in the immune system across life-history stages. *Am Nat* 172:E99–E112. doi:10.1086/589521
- Lummaa V, Clutton-Brock T (2002) Early development, survival and reproduction in humans. *Trends Ecol Evol* 17:141–147. doi:10.1016/S0169-5347(01)02414-4
- Martins TLF (2004) Sex-specific growth rates in zebra finch nestlings: a possible mechanism for sex ratio adjustment. *Behav Ecol* 15:174–180. doi:10.1093/beheco/arg094
- Matson KD, Ricklefs RE, Klasing KC (2005) A hemolysis-hemagglutination assay for characterizing constitutive innate humoral immunity in wild and domestic birds. *Dev Comp Immunol* 29:275–286. doi:10.1016/j.dci.2004.07.006
- Mauck RA, Matson KD, Philipsborn J, Ricklefs RE (2005) Increase in the constitutive innate humoral immune system in Leach's storm-petrel (*Oceanodroma leucorhoa*) chicks is negatively correlated with growth rate. *Funct Ecol* 19:1001–1007. doi:10.1111/j.1365-2435.2005.01060.x
- McGraw KJ, Ardia DR (2005) Sex differences in carotenoid status and immune performance in zebra finches. *Evol Ecol Res* 7:251–262
- Mendes L, Piersma T, Hasselquist D, Matson KD, Ricklefs RE (2006) Variation in the innate and acquired arms of the immune system among five shorebird species. *J Exp Biol* 209:284–291. doi:10.1242/jeb.02015
- Metcalfe NB, Monaghan P (2001) Compensation for a bad start: grow now, pay later? *Trends Ecol Evol* 16:254–260. doi:10.1016/S0169-5347(01)02124-3
- Møller AP, Haussy C (2007) Fitness consequences of variation in natural antibodies and complement in the barn swallow *Hirundo rustica*. *Funct Ecol* 21:363–371. doi:10.1111/j.1365-2435.2006.01215.x
- Møller AP, Saino N (2004) Immune response and survival. *Oikos* 104:299–304. doi:10.1111/j.0030-1299.2004.12844.x
- Monaghan P (2008) Early growth conditions, phenotypic development and environmental change. *Philos Trans R Soc B Biol Sci* 363:1635–1645. doi:10.1098/rstb.2007.0011
- Naguib M, Nemitz A (2007) Living with the past: nutritional stress in juvenile males has immediate effects on their plumage ornaments and on adult attractiveness in zebra finches. *PLoS One* 2:5. doi:10.1371/journal.pone.0000901
- Naguib M, Riebel K, Marzal A, Gil D (2004) Nestling immunocompetence and testosterone covary with brood size in a songbird. *Proc R Soc Lond B Biol Sci* 271:833–838. doi:10.1098/rspb.2003.2673
- Norris K, Evans MR (2000) Ecological immunology: life history trade-offs and immune defense in birds. *Behav Ecol* 11:19–26. doi:10.1093/beheco/11.1.19
- Ochsenbein AF, Zinkernagel RM (2000) Natural antibodies and complement link innate and acquired immunity. *Immunol Today* 21:624–630. doi:10.1016/S0167-5699(00)01754-0
- Parejo D, Silva N, Aviles JM (2007) Within-brood size differences affect innate and acquired immunity in roller *Coracias garrulus* nestlings. *J Avian Biol* 38:717–725. doi:10.1111/j.2007.0908-8857.04081.x

- Roberts ML, Buchanan KL, Evans MR (2004) Testing the immunocompetence handicap hypothesis: a review of the evidence. *Anim Behav* 68:227–239. doi:10.1016/j.anbehav.2004.05.001
- Saino N, Calza S, Møller AP (1997) Immunocompetence of nestling barn swallows in relation to brood size and parental effort. *J Anim Ecol* 66:827–836
- Saino N, Suffritti C, Martinelli R, Rubolini D, Møller AP (2003) Immune response covaries with corticosterone plasma levels under experimentally stressful conditions in nestling barn swallows (*Hirundo rustica*). *Behav Ecol* 14:318–325. doi:10.1093/beheco/14.3.318
- Sheldon BC, Verhulst S (1996) Ecological immunology: costly parasite defences and trade-offs in evolutionary ecology. *Trends Ecol Evol* 11:317–321
- Soler JJ, Martin-Vivaldi M, Haussy C, Møller AP (2007) Intra- and interspecific relationships between nest size and immunity. *Behav Ecol* 18:781–791. doi:10.1093/beheco/arm045
- Stier KS, Almasi B, Gasparini J, Piau R, Roulin A, Jenni L (2009) Effects of corticosterone on innate and humoral immune functions and oxidative stress in barn owl nestlings. *J Exp Biol* 212:2084–2090. doi:10.1242/jeb.024406
- Talloe W, Lens L, Van Dongen S, Adriaensen F, Matthysen E (2010) Mild stress during development affects the phenotype of great tit *Parus major* nestlings: a challenge experiment. *Biol J Linn Soc* 100:103–110. doi:10.1111/j.1095-8312.2010.01422.x
- Tschirren B, Rutstein AN, Postma E, Mariette M, Griffith SC (2009) Short- and long-term consequences of early developmental conditions: a case study on wild and domesticated zebra finches. *J Evol Biol* 22:387–395. doi:10.1111/j.1420-9101.2008.01656.x
- Verhulst S, Holveck MJ, Riebel K (2006) Long-term effects of manipulated natal brood size on metabolic rate in zebra finches. *Biol Lett* 2:478–480. doi:10.1098/rsbl.2006.0496
- Wiersma P, Verhulst S (2005) Effects of intake rate on energy expenditure, somatic repair and reproduction of zebra finches. *J Exp Biol* 208:4091–4098. doi:10.1242/jeb.01854
- Wiersma P, Salomons HM, Verhulst S (2005) Metabolic adjustments to increasing foraging costs of starlings in a closed economy. *J Exp Biol* 208:4099–4108. doi:10.1242/jeb.01855
- Zann RA (1996) *The zebra finch: a synthesis of field and laboratory studies*. Oxford University Press, Oxford